

Patient Care Cervical Cancer Screening

i4C Indicator PHC-CAN-003: Patient Care Cervical Cancer Screening v 2.1

Category: Preventive Health Care/Cancer

Purpose:

This indicator is used to assist with the management of preventive screening for cervical cancer.

Source:

Based on the HQO indicator framework version published in October 2015.

Base Population:

All female patients age 21 to 69 inclusive, with an Active demographic status recorded in the EMR.

Indicator Segments:

Screening up to date: Count of female patients age 21-69 WITHOUT a pap exclusion documented AND WITH a pap screening documented in the past 36 months.

Screening overdue: Count of female patients age 21-69 WITHOUT a pap exclusion documented AND WITHOUT a pap screening documented in the past 36 months.

Excluded: Count of female patients age 21-69 WITH a pap exclusion documented.

Pap Exclusion can be documented as:

- A Q140A service code that has been submitted for a patient
- ICD-9 diagnosis code V45.77 (Acquired absence of organ, genital organs) entered as a current or past diagnosis in the cumulative patient profile
- Any of the following ICD-9 procedure codes entered as a procedure in past medical/surgical history of the cumulative patient profile:
 - 68.3 (Subtotal Abdominal Hysterectomy)
 - 68.4 (Total Abdominal Hysterectomy)
 - 68.5 (Vaginal Hysterectomy)
 - 68.6 (Radical Abdominal Hysterectomy)
 - 68.7 (Radical Vaginal Hysterectomy)
 - 68.8 (Pelvic evisceration)
 - 68.9 (Other and unspecified hysterectomy)
- Any of the following text documented as a current/past diagnosis or as a procedure within past medical/surgical history of the cumulative patient profile:
 - 'Cervical Ca'
 - 'Hysterectomy'
 - 'Hysterosal'
 - 'TVH'
 - 'TABH'
 - 'TAH'

Pap Screening can be documented as a Q011A tracking code that has been submitted for a patient, a report from a sending facility identifying that a pap test has been completed, or as a procedure documented within Past Medical/Surgical history in the cumulative patient profile.

Suggested Indicator Use:

Physician or practice use of this indicator is to review charts of patients overdue for a Pap test in order to manage the recall of patients.