PAS-II

HAQ-II

1. Place an X in the box which best describes your usual abilities *over the past week.*

Are you able to:	Without **ANY** Difficulty	With **SOME** Difficulty	With **MUCH** Difficulty	**UNABLE** To Do
a. Get on and off the toilet?	_____0	_____1	_____2	_____3
b. Walk outdoors on flat ground?	_____0	_____1	_____2	_____3
c. Wait in a line for 15 minutes?	_____0	_____1	_____2	_____3
d. Reach and get down a 5-pound object (such as a bag of sugar) from just above your head?	_____0	_____1	_____2	_____3
e. Do outside work (such as yard work)?	_____0	_____1	_____2	_____3
f. Go up 2 or more flights of stairs?	_____0	_____1	_____2	_____3
g. Move heavy objects?	_____0	_____1	_____2	_____3
h. Stand up from a straight chair?	_____0	_____1	_____2	_____3
i. Lift heavy objects?	_____0	_____1	_____2	_____3
j. Open car doors?	_____0	_____1	_____2	_____3

Pain Scale

2. How much pain have you had because of your condition OVER THE PAST WEEK? Please indicate below how severe your pain has been:

NO PAIN ☐ SEVERE PAIN

0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5 7.0 7.5 8.0 8.5 9.0 9.5 10

Patient global assessment

3. Considering all the ways in which illness and health may affect you at this time, please indicate below how you are doing:

VERY WELL ☐ VERY POORLY

0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5 7.0 7.5 8.0 8.5 9.0 9.5 10

FOR PHYSICIAN USE ONLY

1. HAQ-II/10

☐ * 3.3

☐

2. Pain Score

+ ☐

3. Patient Global

+ ☐

Raw Score

☐ /3

Conversion table (Raw Score /3):

1=0.3 16=5.3
2=0.7 17=5.7
3=1.0 18=6.0
4=1.3 19=6.3
5=1.7 20=6.7
6=2.0 21=7.0
7=2.3 22=7.3
8=2.7 23=7.7
9=3.0 24=8.0
10=3.3 25=8.3
11=3.7 26=8.7
12=4.0 27=9.0
13=4.3 28=9.3
14=4.7 29=9.7
15=5.0 30=10

PAS-II

☐

Interpretation

Remission: ≤0.25
Minimal: ≤3.7
Moderate: <8.0
Severe: ≥8.0

Adapted from
1. Wolfe F, Michaud K, Pincus T. A composite disease activity scale for clinical practice, observational studies, and clinical trials: The patient activity scale (PAS/PAS-II). J Rheumatol. 2005 Dec;32(12):2410-5.