References

1. Rheumatoid arthritis: Physician performance measurement set [homepage on the Internet]. American Medical Association and National Committee for Quality Assurance. 2008 7/2008. Available from: http://www.ama-assn.org/ama1/pub/upload/mm/370/rheumatoid-arthritis.pdf.

2. Saag KG, Teng GG, Patkar NM, Anuntiyo J, Finney C, Curtis JR, et al. American college of rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. Arthritis Rheum. 2008 Jun 15;59(6):762-84.

3. Boers M, Brooks P, Strand CV, Tugwell P. The OMERACT filter for outcome measures in rheumatology. J Rheumatol. 1998 Feb;25(2):198-9.

4. Fransen J, Stucki G, van Riel PLCM. Rheumatoid arthritis measures: Disease activity score (DAS), disease activity score-28 (DAS28), rapid assessment of disease activity in rheumatology (RADAR), and rheumatoid arthritis disease activity index (RADAI). AC&R. 2003;49(S5):S214-24.

5. Aletaha D, Smolen J. The simplified disease activity index (SDAI) and the clinical disease activity index (CDAI): A review of their usefulness and validity in rheumatoid arthritis. Clin Exp Rheumatol. 2005 Sep-Oct;23(5 Suppl 39):S100-8.

6. Smolen JS, Breedveld FC, Schiff MH, Kalden JR, Emery P, Eberl G, et al. A simplified disease activity index for rheumatoid arthritis for use in clinical practice. Rheumatology (Oxford). 2003 Feb;42(2):244-57.

7. Aletaha D, Smolen J, Ward MM. Measuring function in rheumatoid arthritis: Identifying reversible and irreversible components. Arthritis Rheum. 2006 Sep;54(9):2784-92.

8. Rintelen B, Haindl PM, Maktari A, Nothnagl T, Hartl E, Leeb BF. SDAI/CDAI levels in rheumatoid arthritis patients are highly dependent on patient's pain perception and gender. Scand J Rheumatol. 2008 Nov-Dec;37(6):410-3.

9. Soubrier M, Zerkak D, Gossec L, Ayral X, Roux C, Dougados M. Which variables best predict change in rheumatoid arthritis therapy in daily clinical practice? J Rheumatol. 2006 Jul;33(7):1243-6.

10. Aletaha D, Smolen JS. Remission of rheumatoid arthritis: Should we care about definitions? Clin Exp Rheumatol. 2006 Nov-Dec;24(6 Suppl 43):S,45-51.

11. Aletaha D, Nell VP, Stamm T, Uffmann M, Pflugbeil S, Machold K, et al. Acute phase reactants add little to composite disease activity indices for rheumatoid arthritis: Validation of a clinical activity score. Arthritis Res Ther. 2005;7(4):R796-806.

12. Wolfe F, Michaud K, Pincus T. A composite disease activity scale for clinical practice, observational studies, and clinical trials: The patient activity scale (PAS/PAS-II). J Rheumatol. 2005 Dec;32(12):2410-5.

13. Wolfe F, Hawley DJ, Cathey MA. Clinical and health status measures over time: Prognosis and outcome assessment in rheumatoid arthritis. J Rheumatol. 1991 Sep;18(9):1290-7.

14. Wolfe F, Michaud K, Pincus T. Development and validation of the health assessment questionnaire II: A revised version of the health assessment questionnaire. Arthritis Rheum. 2004 Oct;50(10):3296-305.

15. Anderson J, Wolfe F, Michaud K. Converting MHAQ, MDHAQ and HAQII scores into HAQ scores using models developed with 33575 RA patients. Arthritis and Rheumatism. 2009;60(S10):S29.

16. Pincus T, Bergman MJ, Yazici Y, Hines P, Raghupathi K, Maclean R. An index of only patient-reported outcome measures, routine assessment of patient index data 3 (RAPID3), in two abatacept clinical trials: Similar results to disease activity score (DAS28) and other RAPID indices that include physician-reported measures. Rheumatology (Oxford). 2008 Mar;47(3):345-9.

17. Pincus T. A multidimensional health assessment questionnaire (MDHAQ) for all patients with rheumatic diseases to complete at all visits in standard clinical care. Bull NYU Hosp Jt Dis. 2007;65(2):150-60.

18. Pincus T, Sokka T. Can a multi-dimensional health assessment questionnaire (MDHAQ) and routine assessment of patient index data (RAPID) scores be informative in patients with all rheumatic diseases? Best Pract Res Clin Rheumatol. 2007 Aug;21(4):733-53.

19. Yazici Y, Bergman M, Pincus T. Time to score quantitative rheumatoid arthritis measures: 28-joint count, disease activity score, health assessment questionnaire (HAQ),

multidimensional HAQ (MDHAQ), and routine assessment of patient index data (RAPID) scores. J Rheumatol. 2008 Apr;35(4):603-9.

20. Pincus T, Yazici Y, Bergman M, Maclean R, Harrington T. A proposed continuous quality improvement approach to assessment and management of patients with rheumatoid arthritis without formal joint counts, based on quantitative routine assessment of patient index data (RAPID) scores on a multidimensional health assessment questionnaire (MDHAQ). Best Pract Res Clin Rheumatol. 2007 Aug;21(4):789-804.

21. Pincus T, Swearingen CJ, Bergman M, Yazici Y. RAPID3 (routine assessment of patient index data 3), a rheumatoid arthritis index without formal joint counts for routine care: Proposed severity categories compared to disease activity score and clinical disease activity index categories. J Rheumatol. 2008 Nov;35(11):2136-47.

22. Theodore Pincus. RAPID3 (routine assessment of patient index data) on an MDHAQ (multidimensional health assessment questionnaire): Agreement with DAS28 (disease activity score) and CDAI (clinical disease activity index) activity categories, scored in five versus more than ninety seconds. Arthritis care And research. 2010;62(2):181.

23. Pincus T, Strand V, Koch G, Amara I, Crawford B, Wolfe F, et al. An index of the three core data set patient questionnaire measures distinguishes efficacy of active treatment from that of placebo as effectively as the american college of rheumatology 20% response criteria (ACR20) or the disease activity score (DAS) in a rheumatoid arthritis clinical trial. Arthritis Rheum. 2003 Mar;48(3):625-30.

24. Scott DL, Choy EH, Greeves A, Isenberg D, Kassinor D, Rankin E, et al. Standardising joint assessment in rheumatoid arthritis. Clin Rheumatol. 1996 Nov;15(6):579-82.

25. Klinkhoff AV, Bellamy N, Bombardier C, Carette S, Chalmers A, Esdaile JM, et al. An experiment in reducing interobserver variability of the examination for joint tenderness. J Rheumatol. 1988 Mar;15(3):492-4.

26. Thompson PW, Hart LE, Goldsmith CH, Spector TD, Bell MJ, Ramsden MF. Comparison of four articular indices for use in clinical trials in rheumatoid arthritis: Patient, order and observer variation. J Rheumatol. 1991 May;18(5):661-5.

27. Pincus T, Callahan LF, Brooks RH, Fuchs HA, Olsen NJ, Kaye JJ. Self-report questionnaire scores in rheumatoid arthritis compared with traditional physical, radiographic, and laboratory measures. Ann Intern Med. 1989 Feb 15;110(4):259-66.

28. Callahan LF, Smith WJ, Pincus T. Self-report questionnaires in five rheumatic diseases: Comparisons of health status constructs and associations with formal education level. Arthritis Care Res. 1989 Dec;2(4):122-31.

29. Rintelen B, Haindl PM, Sautner J, Leeb BA, Deutsch C, Leeb BF. The rheumatoid arthritis disease activity index-5 in daily use. proposal for disease activity categories. J Rheumatol. 2009 May;36(5):918-24.

30. Leeb BF, Haindl PM, Maktari A, Nothnagl T, Rintelen B. Patient-centered rheumatoid arthritis disease activity assessment by a modified RADAI. J Rheumatol. 2008 Jul;35(7):1294-9.

31. Rintelen B, Sautner J, Haindl P, Andel I, Maktari A, Leeb B. Comparison of three rheumatoid arthritis disease activity scores in clinical routine. Scand J Rheumatol. 2009 Jul 7:1-7.

32. Leeb BF, Sautner J, Mai HT, Haindl PM, Deutsch C, Rintelen B. A comparison of patient questionnaires and composite indexes in routine care of rheumatoid arthritis patients. Joint Bone Spine. 2009 Jun 9.

33. Salaffi F, Peroni M, Ferraccioli GF. Discriminating ability of composite indices for measuring disease activity in rheumatoid arthritis: A comparison of the chronic arthritis systemic index, disease activity score and thompson's articular index. Rheumatology (Oxford). 2000 Jan;39(1):90-6.

34. Ferraccioli GF, Salaffi F, Troise-Rioda W, Bartoli E. The chronic arthritis systemic index (CASI). Clin Exp Rheumatol. 1994 May-Jun;12(3):241-7.

35. Ferraccioli G, Bartoli E, Salaffi F, Peroni M. The chronic arthritis systemic index: A nomogram to assess the activity and severity of chronic arthritis. Arthritis Rheum. 1993 Aug;36(8):1180-1.

36. Pincus T, Bergman M, Sokka T, Roth J, Swearingen C, Yazici Y. Visual analog scales in formats other than a 10 centimeter horizontal line to assess pain and other clinical data. J Rheumatol. 2008 Aug;35(8):1550-8.

37. Scott PJ, Huskisson EC. Measurement of functional capacity with visual analogue scales. Rheumatol Rehabil. 1977 Nov;16(4):257-9.

38. Wells GA, Boers M, Shea B, Brooks PM, Simon LS, Strand CV, et al. Minimal disease activity for rheumatoid arthritis: A preliminary definition. J Rheumatol. 2005 Oct;32(10):2016-24.

39. Rohekar G, Pope J. Test-retest reliability of patient global assessment and physician global assessment in rheumatoid arthritis. J Rheumatol. 2009 Sep 15.

40. Lassere MN, van der Heijde D, Johnson KR, Boers M, Edmonds J. Reliability of measures of disease activity and disease damage in rheumatoid arthritis: Implications for smallest detectable difference, minimal clinically important difference, and analysis of treatment effects in randomized controlled trials. J Rheumatol. 2001 Apr;28(4):892-903.

41. Hernandez-Cruz B, Cardiel MH. Intra-observer reliability of commonly used outcome measures in rheumatoid arthritis. Clin Exp Rheumatol. 1998 Jul-Aug;16(4):459-62.

42. Nicolau G, Yogui MM, Vallochi TL, Gianini RJ, Laurindo IM, Novaes GS. Sources of discrepancy in patient and physician global assessments of rheumatoid arthritis disease activity. J Rheumatol. 2004 Jul;31(7):1293-6.

43. Aletaha D, Machold KP, Nell VP, Smolen JS. The perception of rheumatoid arthritis core set measures by rheumatologists. results of a survey. Rheumatology (Oxford). 2006 Sep;45(9):1133-9.

44. Harrington JT. The uses of disease activity scoring and the physician global assessment of disease activity for managing rheumatoid arthritis in rheumatology practice. J Rheumatol. 2009 May;36(5):925-9.

45. Wolfe F, Michaud K, Pincus T, Furst D, Keystone E. The disease activity score is not suitable as the sole criterion for initiation and evaluation of anti-tumor necrosis factor therapy in the clinic: Discordance between assessment measures and limitations in questionnaire use for regulatory purposes. Arthritis Rheum. 2005 Dec;52(12):3873-9.

46. Hanly JG, Mosher D, Sutton E, Weerasinghe S, Theriault D. Self-assessment of disease activity by patients with rheumatoid arthritis. J Rheumatol. 1996 Sep;23(9):1531-8.

47. Strand V, Cohen S, Crawford B, Smolen JS, Scott DL, Leflunomide Investigators Groups. Patient-reported outcomes better discriminate active treatment from placebo in randomized controlled trials in rheumatoid arthritis. Rheumatology (Oxford). 2004 May;43(5):640-7.

48. van Riel PL, Freundlich B, MacPeek D, Pedersen R, Foehl JR, Singh A, et al. Patient-reported health outcomes in a trial of etanercept monotherapy versus combination therapy with etanercept and methotrexate for rheumatoid arthritis: The ADORE trial. Ann Rheum Dis. 2008 Aug;67(8):1104-10.

49. Scott J, Huskisson EC. Vertical or horizontal visual analogue scales. Ann Rheum Dis. 1979 Dec;38(6):560.

50. Athale N, Sturley A, Skoczen S, Kavanaugh A, Lenert L. A web-compatible instrument for measuring self-reported disease activity in arthritis. J Rheumatol. 2004 Feb;31(2):223-8.

51. Yazici Y, Erkan D, Peterson MG, Kagen LJ. Morning stiffness: How common is it and does it correlate with physician and patient global assessment of disease activity? J Rheumatol. 2001 Jun;28(6):1468-9.

52. Baker PR, Groh JD, Kraag GR, Tugwell P, Wells GA, Boisvert D. Impact of patient with patient interaction on perceived rheumatoid arthritis overall disease status. Scand J Rheumatol. 1996;25(4):207-12.

53. Ward MM. Clinical measures in rheumatoid arthritis: Which are most useful in assessing patients? J Rheumatol. 1994 Jan;21(1):17-27.

54. Berglin E, Lorentzon R, Nordmark L, Nilsson-Sojka B, Rantapaa Dahlqvist S. Predictors of radiological progression and changes in hand bone density in early rheumatoid arthritis. Rheumatology (Oxford). 2003 Feb;42(2):268-75.

55. B F Leeb. Lack of agreement between patients and physicians perspectives of rheumatoid arthritis disease activity changes. Scandinavian Journal of Rheumatology. 2006;35; 35(6; 6):441; 441~!~.

56. van der Heijde DM, van't Hof MA, van Riel PL, van Leeuwen MA, van Rijswijk MH, van de Putte LB. Validity of single variables and composite indices for measuring disease activity in rheumatoid arthritis. Ann Rheum Dis. 1992 Feb;51(2):177-81.

57. Stucki G, Liang MH, Stucki S, Bruhlmann P, Michel BA. A self-administered rheumatoid arthritis disease activity index (RADAI) for epidemiologic research. psychometric properties and correlation with parameters of disease activity. Arthritis Rheum. 1995 Jun;38(6):795-8.

58. Wolfe F, Kleinheksel SM, Cathey MA, Hawley DJ, Spitz PW, Fries JF. The clinical value of the stanford health assessment questionnaire functional disability index in patients with rheumatoid arthritis. J Rheumatol. 1988 Oct;15(10):1480-8.