Patient global assessment

Considering all the ways in which illness and health may affect you at this time, please indicate below how you are doing:

VERY WELL ☐ VERY POORLY
0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5 7.0 7.5 8.0 8.5 9.0 9.5 10

Considering all the ways in which illness and health may affect you at this time, please indicate below how you are doing:

VERY WELL O O O O O O O O O O O O O O O O O O O O O **VERY POORLY**
0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5 7.0 7.5 8.0 8.5 9.0 9.5 10

Considering all the ways in which illness and health may affect you at this time, please indicate below how you are doing:

VERY WELL |————————————————————————| **VERY POORLY**

Proposed definitions of low disease activity are PtGA ≤ 12.5(24) and ≤ 20.(25)

Adapted from

(1) Wells GA, Boers M, Shea B, Brooks PM, Simon LS, Strand CV, et al. Minimal disease activity for rheumatoid arthritis: A preliminary definition. J Rheumatol. 2005 Oct;32(10):2016-24.
(2) Scott PJ, Huskisson EC. Measurement of functional capacity with visual analogue scales. Rheumatol Rehabil. 1977 Nov;16(4):257-9.
(3) Pincus T, Bergman M, Sokka T, Roth J, Swearingen C, Yazici Y. Visual analog scales in formats other than a 10 centimeter horizontal line to assess pain and other clinical data. J Rheumatol. 2008 Aug;35(8):1550-8.