# Provider global assessment




**Provider assessment of global disease activity:**

**NO ACTIVITY** O O O O O O O O O O O O O O O O O O O O O **MAXIMUM ACTIVITY**

0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5 7.0 7.5 8.0 8.5 9.0 9.5 10

**Provider assessment of global disease activity:**

**NO ACTIVITY** |————————————————————————| **MAXIMUM ACTIVITY**

**Interpretation**

- A proposed definition of low disease activity is PrGA ≤ 15.
- Grouping disease activity into no disease activity (0), mild (>0-3.0), moderate (>3.0-6.5), and severe (>6.5-10) has also been suggested.

Adapted from


(1) Wells GA, Boers M, Shea B, Brooks PM, Simon LS, Strand CV, et al. Minimal disease activity for rheumatoid arthritis: A preliminary definition. J Rheumatol. 2005 Oct;32(10):2016-24.

(2) Scott PJ, Huskisson EC. Measurement of functional capacity with visual analogue scales. Rheumatol Rehabil. 1977 Nov;16(4):257-9.

(3) Pincus T, Bergman M, Sokka T, Roth J, Swearingen C, Yazici Y. Visual analog scales in formats other than a 10 centimeter horizontal line to assess pain and other clinical data. J Rheumatol. 2008 Aug;35(8):1550-8.