

COPD Smoking Counselling

I4C Indicator CDM-COPD-002: COPD Smoking Counselling v 2.0

Category: CDM/COPD

Purpose:

This indicator is used to identify patients with COPD who may be eligible for smoking cessation counselling.

Source:

Based on the HQO Quality Standards for COPD Published 2018.

Base Population:

All patients with an Active demographic status recorded in the EMR who have a confirmed COPD diagnosis through entry of an appropriate diagnosis code.

Indicator Segments:

Smokers With Counselling: Count of COPD patients documented as current smokers in the EMR with one or more smoking cessation billing codes (E079 or K039) submitted within the past 12 months.

Smokers Without Counselling: Count of COPD patients documented as current smokers in the EMR without any smoking cessation billing codes (E079 or K039) submitted within the past 12 months.

Non-smokers: Count of COPD patients documented as Non-smokers (including never smoked or former smoker) in the EMR.

Smoking Not Recorded: Count of COPD patients documented in the EMR who have never had smoking status documented.

For a complete list of diagnosis codes used to identify COPD, see COPD Identification (CDM-COPD-001).

Suggested Indicator Use:

Physician or practice use of this indicator is to identify COPD patients to whom smoking cessation support can be offered and patients who have been offered smoking cessation for whom smoking cessation billing codes can be submitted. This indicator can also be used to identify COPD patients without smoking status documented.